



## Source

Anti-SARS-CoV-2 Spike RBD Antibody, Chimeric mAb, Human IgG1 (S1N-M130) is a chimeric monoclonal antibody combining the constant domains of the human IgG1 molecule with mouse variable regions. The variable region was obtained from a mouse immunized with purified recombinant SARS-CoV-2 Spike S1 Protein. *This antibody can broadly recognize all Variants of Concerns (VOCs), including Alpha (B.1.1.7), Beta (B.1.351), Gamma (P.1), Delta (B.1.617.2) and Omicron (B.1.1.529).*

## Isotype

Human IgG1 | Human Kappa

## Conjugate

Unconjugated

## Reactivity

Virus

## Specificity

This product is a specific antibody against SARS-CoV-2 Spike protein RBD domain. No cross-reactivity is detected with Spike protein RBD domain of other coronaviruses, including SARS-CoV, MERS-CoV, HCoV-229E, HCoV-NL63, HCoV-OC43 and HCoV-HKU1.

## Application

Application	Recommended Usage
ELISA	0.2-50 ng/mL

## Purity

>95% as determined by SDS-PAGE.

## Purification

Protein A purified/ Protein G purified

## Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

## Reconstitution

Please see Certificate of Analysis for specific instructions.

*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*

## Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

*Please avoid repeated freeze-thaw cycles.*

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

## SDS-PAGE



Anti-SARS-CoV-2 Spike RBD Antibody, Chimeric mAb, Human IgG1 on SDS-PAGE under reducing (R) and non-reducing (NR) conditions. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.


## Bioactivity-ELISA

Discounts, Gifts,  
and more!






Anti-SARS-CoV-2 Spike RBD Antibody, Chimeric mAb, Human IgG1 ELISA  
0.2 µg of SARS-CoV-2 S1 protein, His Tag per well




Immobilized SARS-CoV-2 S1 protein, His Tag (Cat. No. S1N-C52H2) at 2 µg/mL (100 µL/well) can bind Anti-SARS-CoV-2 Spike RBD Antibody, Chimeric mAb, Human IgG1 (Cat. No. S1N-M130) with a linear range of 0.2-3 ng/mL (QC tested).

Anti-SARS-CoV-2 Spike RBD Antibody, Chimeric mAb, Human IgG1 (S1N-M130) bind to SARS-CoV-2 Spike Trimer, His Tag (B.1.1.529/Omicron) ELISA




Immobilized Anti-SARS-CoV-2 Spike RBD Antibody, Chimeric mAb, Human IgG1 (S1N-M130) at 1 µg/mL (100 µL/well) can bind SARS-CoV-2 Spike Trimer, His Tag (B.1.1.529/Omicron) (Cat.No. SPN-C52Hz) with a linear range of 0.04-0.125 µg/mL (Routinely tested).

Anti-SARS-CoV-2 Spike RBD Antibody, Chimeric mAb, Human IgG1 (S1N-M130) bind to SARS-CoV-2 Spike RBD, His Tag (B.1.1.529/Omicron) ELISA




Immobilized Anti-SARS-CoV-2 Spike RBD Antibody, Chimeric mAb, Human IgG1 (S1N-M130) at 1 µg/mL (100 µL/well) can bind SARS-CoV-2 Spike RBD, His Tag (B.1.1.529/Omicron) (Cat.No. SPD-C522e) with a linear range of 0.0005-0.004 µg/mL (Routinely tested).

## Bioactivity-BLI




Loaded Anti-SARS-CoV-2 Spike RBD Antibody, Chimeric mAb, Human IgG1 (Cat. No. S1N-M130) on AHC Biosensor, can bind SARS-CoV-2 S protein RBD, His Tag (Cat. No. SPD-C52H3) with an affinity constant of 0.603 nM as determined in BLI assay (ForteBio Octet Red96e) (Routinely tested).




Loaded Anti-SARS-CoV-2 Spike RBD Antibody, Chimeric mAb, Human IgG1 (Cat. No. S1N-M130) on AHC Biosensor, can bind SARS-CoV-2 S protein RBD (N501Y), His Tag (Cat. No. SPD-C52Hn) with an affinity constant of 1.38 nM as determined in BLI assay (ForteBio Octet Red96e) (Routinely tested).

Discounts, Gifts,  
and more!







Loaded Anti-SARS-CoV-2 Spike RBD Antibody, Chimeric mAb, Human IgG1 (Cat. No. S1N-M130) on AHC Biosensor, can bind SARS-CoV-2 S protein RBD (K417N, E484K, N501Y), His Tag (Cat. No. SPD-C52Hp) with an affinity constant of 1.10 nM as determined in BLI assay (ForteBio Octet Red96e) (Routinely tested).



Loaded Anti-SARS-CoV-2 Spike RBD Antibody, Chimeric mAb, Human IgG1 (Cat. No. S1N-M130) on AHC Biosensor, can bind SARS-CoV-2 S protein RBD (K417T, E484K, N501Y), His Tag (Cat. No. SPD-C52Hr) with an affinity constant of 1.07 nM as determined in BLI assay (ForteBio Octet Red96e) (Routinely tested).



Loaded Anti-SARS-CoV-2 Spike RBD Antibody, Chimeric mAb, Human IgG1 (Cat. No. S1N-M130) on AHC Biosensor, can bind SARS-CoV-2 Spike RBD (L452R, T478K), His Tag (Cat. No. SPD-C52Hh) with an affinity constant of 1.03 nM as determined in BLI assay (ForteBio Octet Red96e) (Routinely tested).



Loaded Anti-SARS-CoV-2 Spike RBD Antibody, Chimeric mAb, Human IgG1 (Cat. No. S1N-M130) on AHC Biosensor, can bind SARS-CoV-2 Spike RBD, His Tag (BA.2/Omicron) (Cat. No. SPD-C522g) with an affinity constant of 1.04 nM as determined in BLI assay (ForteBio Octet Red96e) (Routinely tested).

### Background

It's been reported that Coronavirus can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity.

### Clinical and Translational Updates

Discounts, Gifts,  
and more!

